

# Selection to outsmart the germs: The evolution of disease

## recognition and social cognition

**Authors:** Sharon E. Kessler<sup>ab\*</sup>, Tyler R. Bonnell<sup>c</sup>, Richard W. Byrne<sup>d</sup>, Colin A. Chapman<sup>a,e</sup>

**Affiliations:** <sup>a</sup>Department of Anthropology, McGill University, <sup>b</sup>Department of Anthropology, Durham University, <sup>c</sup>Department of Psychology, University of Lethbridge, <sup>d</sup>School of Psychology and Neuroscience, University of St. Andrews, <sup>e</sup>Wildlife Conservation Society, New York

### \*Corresponding author:

Sharon E. Kessler  
Department of Anthropology  
McGill University  
855 Sherbrooke West  
Montreal, Quebec H3A 2T7  
Canada  
[Sharon.Kessler@mail.mcgill.ca](mailto:Sharon.Kessler@mail.mcgill.ca)

**Abstract**

The emergence of providing care to diseased conspecifics must have been a turning point during the evolution of hominin sociality. On a population level, such care may have minimized the costs of socially transmitted diseases at a time of increasing social complexity, although individual care-givers would have potentially incurred increased transmission risks while providing care. We propose that care-giving likely originated within kin networks where the costs of providing care may have been balanced by fitness increases obtained through caring for ill kin. We test a novel theory of hominin cognitive evolution in which disease may have selected for the cognitive ability to recognize when a conspecific is infected. Moreover, because diseases may produce symptoms that are likely detectable via the perceptual-cognitive pathways integral to social cognition, we suggest that disease recognition and social cognition may have evolved together. We use agent-based modeling to test 1) under what conditions disease can select for increasing disease recognition and care-giving among kin, 2) whether the strength of selection varies according to the disease's characteristics, 3) whether providing care produces greater selection for cognition than an avoidance strategy, and 4) whether care-giving alters the progression of the disease through the population. We compare the selection created by diseases with different fatality rates (i.e., similar to Ebola, Crimean-Congo hemorrhagic fever, measles, and scabies) under conditions where agents provide care to kin and under conditions where they avoid infected kin. The greatest selection was produced by the measles-like disease which had lower risks to the care-giver and a prevalence that was low enough that it did not disrupt the population's kin networks. When care-giving and avoidance strategies were compared, we found that care-giving reduced the severity of the disease outbreaks and subsequent population crashes. The greatest selection for increased cognitive abilities occurred early in the model runs when the outbreaks and population crashes were most severe. Therefore, we conclude that over the

course of human evolution, repeated introductions of novel diseases into naïve populations could have produced sustained selection for increased disease recognition and care-giving behavior, leading to the evolution of increased cognition, social complexity, and, eventually, medical care in humans. Finally, we lay out predictions derived from our disease recognition hypothesis of hominin cognitive evolution that we encourage paleoanthropologists, bioarchaeologists, primatologists, and paleogeneticists to test.

**Key words:** agent-based model, disease transmission, cooperation, hominin evolution, social complexity, kin selection

## Introduction

Exposure to disease is a major cost of sociality (McCabe et al. 2015; Nunn and Altizer 2006; Rifkin et al. 2012). Despite this, hominins have evolved extraordinary social complexity (Tomasello 2014), including a strikingly social way of mitigating the effects of socially transmitted diseases—we provide care to diseased individuals. Such care hinges on the ability to recognize disease in others. Currently, the cognitive basis of this ability is not well understood. In this paper, we present the novel hypothesis that the ability to recognize disease may have evolved together with social cognition in hominins.

A synthesis of paleoanthropological, ethnographic, and host-parasite research suggests that increasing social complexity during the origin of *Homo* dramatically increased disease risk, i.e., (Harper and Armelagos 2013; McCabe et al. 2015; Rifkin et al. 2012; Sugiyama 2004). Thus, part of the selection for increasing cognitive abilities in *Homo* may have been selection to accurately assess the disease risk presented by interaction partners. We integrate findings from the literature on hominin social structure, hominin disease ecology, disease recognition in nonhuman animals, and human social cognition. Based on these data, we create an agent-based model to examine under what conditions increased cognition and care-giving could have evolved in the hominin lineage. Using our results, we create predictions deriving from our novel disease recognition hypothesis of hominin cognitive evolution that can be tested by paleoanthropologists, paleogeneticists, bioarchaeologists, and primatologists.

## Broadening social networks between hominin subgroups

Across birds and mammals, larger communities show greater levels of contagious parasites, environmentally transmitted parasites, and vector-borne parasites (Rifkin et al. 2012). Though network modularity (sub-grouping) may reduce the transmission risks in large communities where many dyads do not interact (Griffin and Nunn 2012), hominin networks

appear to have connected spatially distant subgroups, facilitating transmission within a fission-fusion, multi-level society (Grove et al. 2012; Hill et al. 2011).

Hominin community sizes have been reconstructed as having expanded over time, from ~50 in apes and small-brained australopiths to 100-120 in late *H. erectus* and *H. heidelbergensis* to 120-150 in *H. neandertalensis* and *H. sapiens* (Aiello and Dunbar 1993; Dunbar 1998; Gamble et al. 2011; Grove et al. 2012; Layton et al. 2012). This is believed to have produced an increase, not only in social network size, but also in complexity (Grove et al. 2012). As hominins dispersed towards northern latitudes and community sizes increased, the home-range requirements for sustaining them would have also increased (Grove et al. 2012). This produced communities whose daily nutritional needs were too large to be fulfilled in the amount of space a cohesive group could cover each day (Grove et al. 2012). The result is thought to have been the evolution of a multi-level fission-fusion system in which larger communities subdivide, rather than foraging cohesively (Grove et al. 2012). This would have enabled large communities of hominins to forage across greater areas and expand into new habitats, yet still obtain the benefits of a large social network, such as information transfer, social learning, and cooperation (Grove et al. 2012; Layton et al. 2012). Thus, even though mean population density decreased over time as hominins dispersed northward, overall community size and social network size likely increased (Grove et al. 2012; Layton et al. 2012).

Community size estimates for modern hunter-gatherers range from 125 to a few thousand (Layton et al. 2012). The extensiveness of human social networks was documented in a study showing that while chimpanzee males typically only interact with about 20 other males, a modern male hunter-gather may watch over 300 other men make tools (Hill et al. 2014). The evolution of such long-distance social networks linking different subgroups (Hill et al. 2014) may have prevented the reduction in disease risk that might otherwise be expected to have

occurred as hominin density decreased, i.e., (Armelagos et al. 2005). Hominins' extensive, community-wide social networks would have facilitated widespread pathogen transmission, including any novel pathogens acquired as hominins spread into new habitats (McCabe et al. 2015).

### **Increasing connectedness within groups**

Simultaneously with the expansion of networks connecting subgroups, the complexity of networks within the subgroups also likely increased with the evolution of cooperative breeding during the origin of *Homo*. Early *Homo* fossil assemblages show an increased number of immature relative to mature individuals compared to australopith assemblages (Tobias 2006), suggesting shortened interbirth intervals, increasing energetic demands on reproducing females, and a shift towards cooperative breeding (Aiello and Key 2002). Ethnographic work supports this view of humans as cooperative breeders, revealing greatly expanded social networks that include multiple providers (hunting males, post-reproductive females) for females and young (Hawkes 2003; Hill et al. 2009; Hrdy 2009). This contrasts with chimpanzees in which the young are solely dependent upon their mothers (Burkart et al. 2009). Collectively, these studies suggest that as community size increased during the origin of *Homo*, so did the complexity of the social networks linking both greater numbers of individuals and different demographics (e.g., young dependents, post-reproductive females, hunting males). The close cooperation, interdependence, and density of social networks within cooperatively breeding hominin groups would have facilitated the spread of diseases within these groups (McCabe et al. 2015).

### **Hominin Disease Ecology**

The shift to larger networks linking subgroups within a larger community and greater connectedness within cooperatively breeding groups is believed to have selected for enhanced social cognition (e.g., prosociality, shared-intentionality, theory of mind) which facilitated prolonged, close interactions among individuals and promoted social learning, cooperation, technological advances and cumulative culture (Burkart et al. 2014; Byrne and Bates 2007; Herrmann et al. 2007; Tomasello et al. 2005; van Schaik et al. 2012; Whiten 2000). However, such intense, close proximity interactions would have also facilitated disease transmission (McCabe et al. 2015). Recent work in genetics and evolutionary medicine indicates that hominins harbored numerous pathogens before the advent of agriculture and animal domestication (Harper and Armelagos 2013). This includes endoparasitic worms (Hoberg et al. 2001; Hurtado et al. 2008), lice (Harper and Armelagos 2013), tuberculosis (Stone et al. 2009), typhoid fever (Harper and Armelagos 2013), whooping cough (Harper and Armelagos 2013), and viruses, e.g., herpes viruses, Epstein Barr virus (Harper and Armelagos 2013). Thus, hominins were likely under strong selection to assess the disease status of others.

#### **Disease recognition in animals and humans**

Comparative evidence suggests that disease recognition may have been present in early hominins (citations below). Several species with relatively low social complexity have been documented to recognize disease, often either avoiding diseased conspecifics or taking advantage of sick and weakened competitors, e.g., social lobsters (Behringer et al. 2006), pipefish (Rosenqvist and Johansson 1995), bullfrog tadpoles (Kiesecker et al. 1999), rodents (Kavaliers et al. 1997), house finches (Bouwman and Hawley 2010; Zylberberg et al. 2012), but see (Nunn 2003). While the underlying cognitive processes are not well understood, these studies suggest that recognition is based on diverse symptoms including olfactory/chemical

cues (Kavaliers et al. 1997; Kiesecker et al. 1999), visual detection of spots (Rosenqvist and Johansson 1995), and behavioral changes including lethargy and feather fluffing (Bouwman and Hawley 2010; Zylberberg et al. 2012). Though the amount of cognitive processing required to detect disease may differ by symptom type, the wide array of cues and recognition in multiple species suggests that some simple form of disease recognition could have been basal in hominins.

Infectious pathogens can cause noticeable symptoms that could potentially be detected via the perceptual-cognitive pathways that are integral to social cognition in primates. Subtle differences perceived in conspecific faces (Leopold and Rhodes 2010; Sartori et al. 2011), voices (Belin 2006; Belin et al. 2004), and movement/gait (Loula et al. 2005; Peterman et al. 2014; Sartori et al. 2011) may enable, not only the decoding of conspecifics' identities, emotions, and intentions, but also facilitate the detection of disease. This could include changes in facial coloration and texture due to fever, rashes, or nasal discharge, changes in vocalizations due to coughing, nasal discharge or reduced lung capacity, and changes in movement/gait due to weakness, lethargy, or signs of pain (Chapman et al. 2005; Fink and Matts 2008; Hart 1988). Thus, if the detection of social information and disease involve the same perceptual-cognitive pathways, then disease circulating within hominin populations may have selected for increased cognitive capacities and care-giving.

Importantly, such disease recognition would *not* require individuals to have an abstract concept of disease. Following the well-accepted definition of cognition as information processing, e.g., seminal book: (Neisser 1967), recent publications: (Byrne and Bates 2007; Deaner et al. 2006; Fernandes et al. 2014; Herrmann et al. 2007; Lee 2007; Reader et al. 2011; Woodley et al. 2015), the cognitive aspect would be processing the proximate cues that distinguish healthy individuals from diseased individuals (changes in appearance, behavior, etc.). Selection for such disease recognition would operate at the ultimate level of



causation (Sherman 1988; Tinbergen 1963), favoring individuals who were able to discriminate who was healthy and who was not. Those who avoided infectious individuals or provided care to ill kin would increase their reproductive fitness. Similarly to how kin recognition can operate without individuals having an abstract concept of kin (Rendall 2004), disease recognition could operate without a concept of disease.

### **Care-giving among animals and humans**

The literature contains numerous reports of striking cases of social care given by animals, including dolphins that cooperatively supported a dying conspecific who could no longer swim (Park et al. 2013), an elephant that attempted to lift a collapsed and dying conspecific to her feet (Douglas-Hamilton et al. 2006), primates that groom, stand watch over, and/or chase others away from dying group members (Anderson et al. 2010; Bezerra et al. 2014; Nakamichi et al. 1996), and an otter group that provisioned an elderly female (Davenport 2010). Though very interesting, these reports do not provide evidence of widespread long-term care which would be expected to have a more significant selective influence on a species' evolution.

Some of the best opportunities for systematically investigating care-giving in animals have come from studies of populations with high prevalences of severe injuries (Beamish and O'Riain 2014; Byrne and Stokes 2002; Stokes and Byrne 2006) or congenital disabilities (Turner et al. 2014). These studies generally suggest that, instead of relying on social care, severely injured or disabled individuals survive by adapting and making adjustments themselves, rather than receiving accommodation or assistance (Beamish and O'Riain 2014; Byrne and Stokes 2002; Stokes and Byrne 2006; Turner et al. 2014). The exception to this is social grooming (Dittus and Ratnayeke 1989). Wound cleaning has been shown to be an important mechanism for avoiding infections and it is widespread in animals (Dittus and

Ratnayeke 1989; Hart 2011). Thus wound cleaning may have been a basal form of social care in hominins.

In addition, evidence from modern foraging, hunting, and horticultural peoples, suggests that provisioning people who are ill or injured is important in reducing the mortality rate (Sugiyama 2004). For example, Sugiyama (2004) found that over 50% of individuals reported at least one time in their lives when they were incapacitated and could not forage for at least a month. During such times, provisioning was critical to their survival (Sugiyama 2004). Based on this evidence, we expect that hominins could have significantly reduced the mortality arising from disease and infection-related injuries through provisioning (Sugiyama 2004) and wound cleaning (Dittus and Ratnayeke 1989). Additionally, food sharing networks of hunting males also served as provisioning networks during times of illness (Gurven et al. 2000; Sugiyama 2004; Sugiyama and Chacon 2000), suggesting that the evolution of social care may have co-evolved with cooperative breeding.

### Care-giving in the fossil record

Fossil evidence of hominins surviving illness, injuries, and disabilities goes back nearly 2 million years to include fossils from *H. erectus*, *H. heidelbergensis*, *H. neandertalensis*, and *H. sapiens*. While the following discussion is not exhaustive, it does illustrate the variety of conditions hominins survived, the time depth of the fossil record, and the taxa included. Below we follow, when possible, the taxonomic classifications provided in Grove et al. (2012). In *H. erectus* this includes: premortem loss of all but one tooth in the 1.77 mya cranium and mandible from Dmanisi (D3444 and D3900 (Lordkipanidze et al. 2005; Lordkipanidze et al. 2006)), possible hypervitaminosis A in the 1.6 mya KNM-ER 1808 (Walker et al. 1982), evidence of a herniated disc in the 1.5-1.6 mya Nariokotome boy KNM-WT 15000 (Grove et al. 2012; Haeusler et al. 2013; Schiess et al. 2014), and a healed cranial

lesion caused by trauma or burning in the 0.6 mya Hulu 1 cranium, also called Nanjing 1 and  
 Tangshan 1 (Shang and Trinkaus 2008; Wu et al. 2011). Among *H. heidelbergensis* this  
 includes craniosynostosis and neurocranial deformities in a 0.53 mya immature, cranium 14,  
 who survived for at least approximately 5 years (Gracia et al. 2009), a 0.53 mya adult male  
 pelvis and lumbar spine, SH Pelvis 1, showing lesions and degeneration possibly resulting  
 from lumbar kyphotic deformity, spondylolisthesis, and Baastrup disease (Bonmati et al.  
 2010), and a squamous temporal lesion that shows healing on the 0.35 mya Broken Hill  
 cranium Kabwe 1 (Grove et al. 2012; McBrearty and Brooks 2000; Montgomery et al. 1994).  
 For Neandertals this includes Aubesier 11, dated to at least 0.17 mya, which shows  
 significant tooth loss and alveolar lesions (Lebel and Trinkaus 2002; Lebel et al. 2001) and  
 Shanidar 1 dated at 73-40 kya who lost much of his right arm, may have been blind on one  
 side, and suffered from hyperostotic disease (Crubezy and Trinkaus 1992; Hublin 2009). *H.*  
*sapiens* individuals that survived severe conditions include: a child, Qafzeh 12 dated to  
 approximate 0.095 mya, who showed signs of hydrocephaly and survived until about 3 years  
 old (Tillier et al. 2001), an older child Qafzeh 11, also dated to 0.95 mya, that had a healed  
 cranial fracture (Coqueugniot et al. 2014), and an adult female, Dolní Věstonice 3, dated to  
 approximately 0.027 mya, who sustained a severe injury to her face that might have  
 interfered with eating (Trinkaus et al. 2006; Trinkaus and Jelinek 1997).

While all of these individuals *might* have benefited from care, comparative evidence with  
 nonhuman primates suggests that care is not necessary (DeGusta 2002, 2003; Dettwyler  
 1991). Studies of wild baboons and great apes show that primates frequently survive even  
 when a hand or foot is maimed or severed, e.g., in snares (Beamish and O'Riain 2014; Byrne  
 and Stokes 2002; Munn 2006; Stokes and Byrne 2006). Though these animals may show  
 changes to their activity budgets (Beamish and O'Riain 2014), altered locomotion patterns  
 (Munn 2006), and reduced feeding efficiency (Byrne and Stokes 2002; Stokes and Byrne

2006), survival appears to be high, with some groups having as many as ~20% of their members permanently disabled (Munn 2006). Extensive tooth loss also appears to be survivable. Apes and other primates have been observed to survive antemortem tooth loss comparable to that observed in the fossil record (Cuozzo and Sauter 2004; DeGusta 2002). Degusta (2002) provides a review of cases in which chimpanzees were observed to survive with tooth loss similar to Aulers 11 and Cuozzo and Sauter (2004) reported that tooth loss is common among ring-tailed lemurs, with one individual surviving with 80% tooth loss. Overall the evidence from the fossil record and animal studies indicate that while various fossils have clearly survived severe health conditions, it is very difficult to rule out the possibility that they may have survived without care (DeGusta 2002, 2003; Dettwyler 1991).

### **The modeling approach**

It is currently not possible to determine when extensive social care evolved in the human lineage, but it is possible to consider *how* it might have evolved and what conditions might have selected for it. We expect that, because kinship is a fundamental property of primate (including human) social networks (Silk 2009), providing care to the diseased may have originated along kin networks. Hamilton's rule of inclusive fitness (Hamilton 1964) predicts that individuals will act altruistically when:  $(\text{benefit to the recipient}) \times (\text{relatedness to recipient}) > (\text{costs to the altruist})$ . Thus, individuals could increase their own reproductive fitness in two ways: 1) by avoiding ill individuals, particularly nonkin, and 2) by providing care to ill kin who, upon recovery, would reproduce. Whether the fitness benefits are greater when individuals avoid ill conspecifics or provide care (thus risking becoming infected) will depend upon the benefits, the degree of relatedness, and the costs.

We use agent-based modeling to test a varying intensity of disease scenarios and quantify selection pressures for increased cognition and care-giving. Agent-based models provide

powerful, quantitative insights into disease transmission, including predicting the impact of current/future outbreaks and planning intervention/prevention strategies, e.g., influenza (Guo et al. 2015), Ebola (Merler et al. 2015). We take the innovative approach of applying these techniques to reconstruct the potential impact of disease on hominin evolution.

A modeling approach is valuable because, while our knowledge is increasing, i.e., (Harper and Armelagos 2013), we do not have sufficiently detailed data concerning how/when disease load changed during hominin evolution to be able to test whether the evolution of care-giving co-occurred with increasing cognitive abilities, social complexity and disease risk. Therefore, we use agent-based modeling to examine under which conditions disease could select for increased cognition and care-giving. We hypothesize that 1) disease will produce care-giving among kin and an increase in average population intelligence, that 2) varying disease characteristics will produce variation in the strength of selection, and that 3) care-giving will produce greater selection for cognition than an avoidance strategy.

## **Material and methods**

### *Study design*

We created two models for comparison. The first (Model 1: Care-giving model) simulates disease transmission in a population of hominins who give care (The ODD description is in Appendix A at the end of the paper. The code is available in supplementary file 1). In order to more fully explore the model and how care-giving may alter the progression of disease through the population, we then created a control model (Model 2: Avoidance only) similar to the first except that agents avoid diseased kin and provide no care. (The ODD description is in Appendix B at the end of the paper. The code is available in supplementary file 2).

### ***Model 1: Care-giving model***

### *Disease characteristics*

We programmed an SIS model (susceptible – infected – susceptible) in Netlogo 5.0.5 (Railsback and Grimm 2011; Wilensky 1999). We created four hypothetical diseases with case fatality rates modeled after Ebola [2014 outbreak: 70% (Aylward et al. 2014; WHO 2014a), Crimean-Congo hemorrhagic fever (40% (WHO 2013), CCHF, hereafter), measles (~10% (WHO 2014b)), and a low risk comparison, such as scabies (fatality rate set at 1%, though scabies is generally not fatal (WHO 2015)]. We did not attempt to precisely simulate the natural history of these diseases. Rather, these diseases were chosen to represent a range of fatality rates occurring in socially transmitted diseases.

### *Optimizing the disease transmission rates*

Because transmission rates have complex relationships with virulence and host density (e.g., trade-off hypothesis (Alizon et al. 2009)), we screened possible transmission rates to determine what would be optimal for persistence of these diseases in this population. For the Ebola-like, CCHF-like, and measles-like diseases, we ran the model 1000 times in Netlogo's BehaviorSpace, varying the probability of transmission from 10-100% by increments of 10. For the scabies-like disease, we ran the model 1000 times varying the probability of transmission from 1% to 98.5% by increments of 2.5. The inclusion of lower transmission values for the scabies-like disease is based on literature showing that less virulent diseases tend propagate slower, e.g., (Alizon et al. 2009; Ewald 1993). Then, for each disease, we selected the runs which had both healthy and diseased individuals after 100 time steps. We averaged the probability of transmission across those successful runs to obtain a transmission rate that is optimal for each respective disease: Ebola-like 78%, CCHF-like 33%, measles-like 10%, scabies-like 2%. The higher transmission rates in the diseases with higher fatality

rates is consistent with the relationship between virulence and transmission documented in the literature (Alizon et al. 2009).

#### *Determining the probability of recovery after care*

We expect that the earliest forms of social care given by hominins would have been assistance with hygiene, including keeping wounds, sores, and topical infections clean as in nonhuman primates (Dittus and Ratnayeke 1989), provisioning those who are too ill to forage with food and water (Sugiyama 2004), and watching over individuals who may be too ill to themselves be vigilant against predators (Anderson et al. 2010; Bezerra et al. 2014; Nakamichi et al. 1996). None of these forms of care requires medical knowledge, yet evidence from nonhuman primates (Dittus and Ratnayeke 1989) and human foraging groups (Sugiyama 2004) suggests that they are effective at reducing mortality rates.

It is difficult to estimate how effective each of these care-giving techniques would be for each of our hypothetical diseases. In nature, the more incapacitated the individual is and the longer the recovery takes, the greater the chances that the individual would succumb to dehydration, starvation, or predation unless care is given. Because we did not wish to bias the effectiveness of the care towards the more severe diseases, we set the probability of recovery after care at 0.5 for all diseases. This reflects an equal chance of recovery and failure to recover.

#### *The population*

The landscape is a 40 x 40 cell grid that wraps horizontally and vertically. Each cell represents 5 km<sup>2</sup>, making the landscape 200 km<sup>2</sup>. This is within the confidence intervals of the space requirements calculated for a community of *H. erectus*, *H. heidelbergensis*, *H. neandertalensis*, and *H. sapiens* using a gas model in Grove et al. (2012). Table 1

summarizes the group sizes, densities, and space requirements presented in Grove et al. (2012).

### [Table 1]

The carrying capacity of the landscape is set at 200. Two hundred was chosen because it is large enough to encompass the group sizes predicted for hominins based on cranial capacities, brain volumes, and neocortex ratios of fossil hominins [Table 1, (Aiello and Dunbar 1993; Gamble et al. 2011; Grove et al. 2012)], but is generally smaller than community sizes reported for modern humans, e.g., (Hill et al. 2014; Layton et al. 2012). We set the carrying capacity above the calculated community sizes for hominins, e.g., ~150 or smaller (Aiello and Dunbar 1993; Dunbar 1998; Gamble et al. 2011; Grove et al. 2012), to allow for the event that the actual community sizes of the model populations would likely be lower than the carrying capacity.

### *Initialization*

The program is initialized with 10 agents randomly placed on the landscape. Each agent is randomly assigned an intelligence score (0-1). In the model the intelligence score is the likelihood of an agent correctly identifying the disease status of another agent. We refer to it as intelligence because we expect that the ability to recognize disease is related to a more general ability for efficient information processing, including social information, e.g., (Byrne and Bates 2007; Deaner et al. 2006; Fernandes et al. 2014; Herrmann et al. 2007; Lee 2007; Reader et al. 2011; Woodley et al. 2015). As the population grows, each offspring's intelligence is drawn from a normal distribution with the parent's intelligence as the mean and a standard deviation of 0.15.

### *Population growth and genetic structure*



The population grows at each time step of the model when healthy agents reproduce according to the formula:  $[(1 - (\text{number of agents} / \text{carrying-capacity})) * \text{number of healthy agents}]$ . Reproduction occurs asexually. Offspring are placed within a radius of 3 of the parent, producing spatial clustering of kin as is consistent with human and nonhuman primate groups (Chapais and Berman 2004; Hatchwell 2010; Hill et al. 2011; Silk 2009).

Relatedness is tracked by links between agents with the links containing the relatedness value. Parent-offspring relationships receive relatedness values of 0.5 and offspring inherit the links of the parent but with  $\frac{1}{2}$  the relatedness value. Because offspring inherit the links of the parent, sibling relationships are included in the model with a relatedness value 0.25. To prevent the model from becoming too computationally intensive, patrilineal relationships and matrilineal relationships beyond a relatedness of 0.25 were not modeled. This decision is supported by findings showing that kin recognition occurs most reliably for close matrilineal kin identified via familiarity, e.g., (Chapais and Berman 2004; Chapais et al. 1997). The population represents a single, kin structured community with multiple matrilineal. Space displays the contact structure between agents and random movement simulates mixing within the population.

### *Space*

With a carrying capacity of 200 individuals and a landscape of 200 km<sup>2</sup>, our model has a maximum population density of 1 individual / km<sup>2</sup>, which is within the confidence intervals calculated for *H. habilis* and *H. erectus* [Table 1, (Grove et al. 2012)]. However, the purpose of our model is not to attempt to reconstruct a particular hominin species or population. We made this decision because the population densities and number of levels of fissioning have been reconstructed to vary dramatically even within species, depending upon the habitat quality and latitude (Atkinson et al. 2008; Grove et al. 2012; Powell et al. 2009). Instead,

hominin societies are conceptualized as more generic fission-fusion communities in which subsets of individuals are out of contact with other subsets of individuals (Grove et al. 2012; Layton et al. 2012). This is represented in our model by the restrictions created by the movement, care-giving, and infection radii. The care-giving radius (5) and infection radius (5) are equal to reflect that agents who are close enough to give care are also close enough to become infected. Similarly, agents who avoid infectious kin by moving away will also be moving away from potential care-givers should they themselves become infected. These radii of 5 represent 25 km<sup>2</sup> and are in the upper range of the distance that modern hunter-gatherers travel from camp when they will return to camp later the same day (Grove et al. 2012; Layton et al. 2012).

#### *Disease and care-giving*

After four time steps of the model, 25 agents are randomly infected with one of the diseases. This is approximately 16% of the population and reliably seeded the disease into the population without increasing to 100% prevalence.

Healthy agents evaluate the relatedness and disease status of other agents within a radius equivalent to 5 grid cells. The infection radius is also set at 5, thus any healthy agent that can provide care, is also close enough to be infected.

Kin are accurately recognized and the accuracy of disease recognition is a function of the agent's intelligence. A random number between 0-1 is drawn. If the number is below the agent's intelligence value, the disease status is correctly recognized. Otherwise, the agent's disease status is incorrectly recognized (healthy kin are classified as diseased or diseased kin are classified as healthy). These individuals make up the group the agent *perceives* to be its diseased kin (perceived diseased kin). Whether the error is a false positive (healthy kin classified as diseased) or a false negative (diseased kin classified as healthy) is determined by

the disease status of the kin agent. Thus, the likelihoods of false positive and false negative errors are functions of disease prevalence. As the proportion of diseased agents increases, false positives decrease and false negatives increase.

Agents randomly select one of their perceived diseased kin and decide whether to provide care based on a modification of Hamilton's rule, which predicts altruism when: (relatedness between the recipient and altruist)\*(benefit to the recipient)>(cost to the altruist) (Hamilton 1964). We adapted this formula so that agents provide care when: (relatedness between the care-giver and the recipient)\*(probability of recovery after care) > (probability of transmission to care-giver)\*(probability of infection being fatal). If the inequality is fulfilled (thus care is given) and the recipient was in fact diseased (not just *perceived* to be diseased), a random number between 0 and 1 is generated and if it is below the probability of recovery, the diseased individual recovers. If the random number was above the probability of recovery, the recipient remains diseased. A new random number is drawn for the care-giver and if it is below the probability of transmission to the care-giver, then the care-giver is infected. If the recipient was erroneously categorized as diseased, but is actually healthy (a false positive error), there is no change in the disease statuses of the recipient or the care-giver. It is worth noting that when a false negative error occurs (diseased kin are classified as healthy), the agent that made the error does not incur a cost that is explicitly coded into the model. However, the agent does potentially incur emergent costs through the interactions between agents. This may occur in two ways: a) if that diseased kin agent dies (later in the model run), this reduces the kin network available to give care, simulating a loss of inclusive fitness to the agent that failed to recognize the disease in its kin, and b) the presence of diseased kin in the population increases the risk that others will become infected, including the agent that failed to recognize the disease in its kin.

If healthy agents have no perceived diseased kin, they move to a grid cell with no other agents on it within a radius of 8. If no empty cells are available, the agent does nothing. A movement radius of 8 represents 40 km<sup>2</sup>. This is the median daily *total* travel distance used by Grove et al. (2012) to calculate hominin area requirements and it is based on data compiled from modern hunter-gathers, e.g., (Layton et al. 2012).

#### *Avoidance of infectious individuals*

If the randomly selected recipient (from the agent's perceived diseased kin) does not fulfill the inequality for receiving care, the agent moves to a grid cell with no other agents on it within a radius of 8. This can occur due to a low relatedness with the recipient of care, high costs of exposure to the disease, or a low likelihood of recipient recovery. Under these conditions, the agent avoids the diseased individual instead of providing care. Note that nonkin do not receive care, thus if no perceived diseased kin are within the care-giving radius, the agent moves.

Because the care-giving radius and the infection radius are set at 5 and this is less than the movement radius (8), agents that do not provide care can move out of the infection radius. The effectiveness of movement as a disease avoidance strategy is based on chance and the density of infected individuals. By chance the healthy agent may move to a grid cell that is outside of the infection radius of the diseased agent. However, as the density of infected agents increases, so does the likelihood that the healthy agent will move to a grid cell that is within the infection radius of another diseased agent. This reflects the difficulties of avoiding exposure at when there is a high density of infectious individuals in the population.

If no empty cells are available, the agent does nothing.

#### *Mortality and disease transmission*

The model generates a random number for each diseased agent. If the number is below the probability of fatality, that agent dies. All healthy agents have a probability of becoming infected from any infected agent within a radius of 5 grid cells, based on the probability of transmission. Five grid cells represent the upper range of the daily travel radius for modern hunter-gatherers (25 km<sup>2</sup>) (Grove et al. 2012; Layton et al. 2012). A random number (0-1) is drawn for each healthy agent in danger of infection. If the number is below the probability of transmission, the agent is infected. If an agent is in danger of infection from more than one diseased agent, the process is repeated for each infectious agent in 5 grid cells.

#### *Model analysis*

We ran the model 2000 times for 100 time steps for each disease. We considered runs to be successfully completed when both the disease and population had persisted (defined as  $\geq 1$  diseased agent and  $\geq 1$  healthy agent at 100 time steps). The first 1000 successfully completed runs were divided into 10 groups of 100. We calculated average population size, average disease prevalence, average percentage of diseased individuals who received care (percent care), and average population intelligence at each time step across the 100 runs. This created an n of 10 average runs for which we made curves depicting the changes in each of these output variables for the four diseases we considered. We used the 10 averages in the subsequent statistical tests instead of the original 1000 runs to avoid inflating our sample size, and thus the power of our tests (Railsback and Grimm 2011).

#### *Statistics*

We compared the endpoints of the curves by comparing the output variables (average population size, average disease prevalence, and average percent care) across the diseases at time step 100 using one-way ANOVAs (n=10 average runs/disease). We calculated the

change in average population intelligence between the first and 100<sup>th</sup> time step, tested whether the differences were different from zero using one-sample T-tests, and whether these differences varied across disease types using a one-way ANOVA. We calculated maximum slopes for the curves of the average percent care and the average population intelligence using `grofit` (Kahm et al. 2010) in R 2.13.1 (RCoreTeam 2011) and RStudio 0.98.1062 (RStudio 2014). We tested whether the slopes differed across disease types using a one-way ANOVA. Some violations of normality and equal variances existed (Supplementary files 3 and 4). One-way t-tests were bootstrapped with 1000 samples for robusticity to non-normality and 95% bias corrected accelerated confidence intervals were calculated (Field 2013). Though one-way ANOVAs are generally robust to such violations when groups have equal sample sizes, when variances were unequal, we used the Brown-Forsythe F-ratio. Alpha was set at 0.05 and multiple comparisons across disease types were Bonferroni corrected when variances were equal and Tamhane T2 corrected when they were unequal. Statistical tests were run in SPSS Statistics 22 or 23 unless otherwise stated.

### ***Model 2: Control model – Avoidance only***

Following the initial analysis of the care-giving model (Model 1), we programmed a control (Model 2: Avoidance only) to further explore how care-giving may have altered the progression of disease through hominin populations. This model used the same population and diseases, but differed in two ways. First, agents who have perceived diseased kin avoid them instead of providing care. All agents with perceived diseased kin move randomly to an empty grid cell within a radius of 8. Second, if the agent has no perceived diseased kin or there are no empty grid cells within a radius of 8, the agent does not move. This differs from the care-giving model in which agents with no diseased kin also move to an empty grid cell within 8. (Because agents that give care do not move, this was necessary in the care-giving

model to ensure movement within the population.) We made this second change to the avoidance model to be conservative with respect to our expectation that only care-giving will produce intelligence changes. This second change increased selection on avoidance behavior because in Model 2 (Avoidance only) the only opportunity agents have to move is when they are avoiding diseased kin.

#### *Model analysis and statistics*

We used the same procedure as above to create 10 average runs for each output variable for each disease. We conducted one-sample T-tests to determine whether the difference in average population intelligence between the first and 100<sup>th</sup> time steps were significantly different from zero for the scabies-like, measles-like, CCHF-like, and Ebola-like diseases. We used two sample T-tests to determine whether the population size, prevalence, and intelligence at the 100<sup>th</sup> time steps differed between models 1 and 2. Some violations of normality and equal variances existed (Supplementary files 3 and 4). T-tests were bootstrapped with 1000 samples for robusticity to non-normality and 95% bias corrected accelerated confidence intervals were calculated (Field 2013). When Levene's test showed violations of the assumptions of equal variances, we report results calculated without assuming equal variances (Field 2013). Alpha was set at 0.05.

#### *Analysis of the intelligence curves produced by Model 1 (Care-giving)*

We analyzed the trajectories of the intelligence curves of the 10 average runs for each disease using linear mixed-models run in R 3.2.4 (RCoreTeam 2015) using the nlme package. We use this approach to relate infection prevalence to changes in mean intelligence, while taking into account population size. We test for an interaction between prevalence and population size on changes to mean intelligence by including interaction term in the model:

prevalence \* population size. As the data are longitudinal (i.e., time series) we allow for autocorrelated errors using an ARMA process, incorporate time as a fixed effect, and use the averaged simulation run as the random effect. We check for issues of multicollinearity using variation inflation factor, and check the residuals of the models for non-normality, heteroscedasticity, and autocorrelation. (model: change in mean intelligence ~ time + prevalence\*population size + random intercept). In order to keep the paper focused on the evolution of increasing average population intelligence, we did not conduct this analysis on the Model 2 curves, which showed either no increase or a decrease in average intelligence.

## Results

### *Model 1: Care-giving model*

After 100 time steps the four diseases produced significantly different population sizes, disease prevalence, percentages of the diseased who received care, and average population intelligences (Tables 2-3, Fig. 1).

**[Table 2]**

**[Table 3]**

**[Figure 1]**

The Ebola-like disease, unlike the other three, produced no care-giving and no change in average population intelligence (Table 4, Fig. 1).

**[Table 4]**



Both the Crimean-Congo hemorrhagic fever-like (CCHF-like) and measles-like diseases show initial increases in both care-giving and intelligence followed by a plateau (Fig. 1). The CCHF-like disease produced a care-giving rate of 4.7%, a final intelligence level of 0.62, and a 12% net change in intelligence. Of the four diseases, the measles-like disease produced the highest rate of care-giving (6.7%) and the highest average population intelligence (0.71) at the final time step. This was generated by the greatest maximum slopes for care-giving and intelligence changes and the greatest net change in intelligence over time (21%). The scabies-like disease showed a strikingly different pattern. As prevalence steadily increased, because the fatality rate was low, care-giving decreased. Infected individuals did not provide care and rarely died, meaning that the number of healthy individuals able to provide care decreased. This produced a negative slope for care-giving, though low increases in average population intelligence were still observed (care-giving rate: 1.4%, final average population intelligence: 0.53, net intelligence change: 3%, Tables 2-3).

#### *Model 2: Control model – Avoidance only*

The model two results revealed two important findings. First, an avoidance strategy did not result in an increase in average population intelligence (Tables 5 and 6). The net change in intelligence overtime was not significantly different from zero under the scabies-like and measles-like conditions (Table 5). Under the CCHF-like and Ebola-like conditions the average population intelligence decreased significantly (Table 5).

#### **[Table 5]**

#### **[Table 6]**

Second, a visual inspection of Figures 2-4 shows that the progression of the diseases through the population differed under Model 1 (care-giving) and Model 2 (avoidance only). Descriptive statistics are provided in Supplementary file 5. For the scabies-like and measles-

like diseases, when agents gave care the final population sizes were higher and the final prevalences were lower (Fig. 2 & 3, Table 6). A visual inspection of Fig. 3b reveals that when agents give care, the “boom and bust” cycle of disease outbreaks in the population was reduced with prevalence increasing and decreasing less dramatically. For the CCHF-like disease, the final population sizes differed however prevalence did not differ (Table 6). An inspection of Fig. 3c shows that the cycle of outbreaks was very similar in the care-giving and avoidance conditions. For the Ebola-like disease, final population size and final prevalence did not differ in the care-giving and avoidance conditions.

[Fig. 2]

[Fig. 3]

[Fig. 4]

#### *Analysis of the intelligence curves produced by Model 1 (care-giving)*

For each of the scabies-like, measles-like, and CCHF-like diseases, time was negatively related to changes in intelligence (Table 7). Thus, the largest increases occurred early in the run with smaller increases occurring later. In the case of the Ebola-like disease, intelligence did not change, thus there was no relationship between time and changes in mean intelligence.

For the scabies-like disease, VIF scores indicated high collinearity between dependent variables (VIF scores >100). When we dropped population size from the analysis, VIF scores fell below 7. In this reduced analysis, changes in intelligence were positively related with prevalence (Table 7, Fig. 5).

For the measles-like disease, changes in intelligence were positively related with both prevalence and population size with the greatest increases in intelligence occurring at larger population sizes and high prevalences (Fig. 6). For the CCHF-like disease, the proportion of

the variation explained by the analysis (marginal  $R^2 = 0.15$ ) was reduced compared to the measles-like (marginal  $R^2 = 0.57$ ) and scabies-like (marginal  $R^2 = 0.47$ ) diseases. However, similar to the measles-like disease, an interaction effect between prevalence and population size was present, indicating that at low prevalences, changes in intelligence were negatively related to population size, but at higher prevalences, they were positively related with population size (Fig. 7). Thus the greatest changes in intelligence occurred at low prevalences and low population sizes or high prevalences and high population sizes.

No relationships between time, prevalence or population size were found for the Ebola-like disease because the Ebola-like disease produced no changes in intelligence (Tables 5 and 7, Fig. 8).

**[Table 7]**

**[Fig. 5]**

**[Fig. 6]**

**[Fig. 7]**

**[Fig. 8]**

## **Discussion**

### *General discussion*

Our findings suggest that the evolution of care-giving may have created a profound shift in how hominins evolved in the presence of their pathogens. The avoidance approach (Model 2) likely represents the basal condition, under which disease either does not select for or against increasing cognitive abilities (high prevalence, low fatality diseases) or selects against it (low prevalence, high fatality diseases). In contrast, under the care-giving condition (Model 1), care-giving not only selected for increasing cognitive abilities, but also altered and

controlled the progression of some of the diseases throughout the population. We discuss both models and their implications in detail below.

### *Model 1*

Our results from Model 1 suggest that disease circulating among kin can select for care-giving among kin and greater cognitive abilities. Furthermore, the diseases produced selection of varying strengths, with higher care-giving rates producing greater increases in average population intelligence.

The findings are relevant to the evolution of care-giving in hominins as they suggest that not all diseases produce care-giving behavior. The high fatality and transmission rates of the Ebola-like disease, when applied to Hamilton's rule (Hamilton 1964), generated costs that were greater than the benefits of care-giving, even to close relatives, thus, all agents avoided ill kin, rather than providing care. Such diseases are not likely to have facilitated the evolution of care-giving or increased social cognition. The CCHF-like disease had intermediate probabilities of fatality and transmission, leading to care-giving only to close kin (parents and offspring:  $r=0.5$ ), and not to more distant relatives like grandparents, grandchildren, or siblings ( $r=0.25$ ) who were avoided when ill. This produced substantial care-giving behavior and selection for increasing intelligence, but the selection was weaker than for the measles-like disease, where care was given to both close and more distant relatives. The scabies-like disease, while it produced care-giving for both close and more distant relatives, produced only low rates of care-giving and correspondingly weak selection for increasing intelligence. These effects result from the very low fatality rate of the scabies-like disease; the population size appears to have been regulated largely by the carrying capacity set in the model (i.e., habitat supports 200 individuals) rather than by the disease. Therefore, as disease prevalence increased, there was a lack of healthy individuals who could

provide care to their diseased kin, leading to a low rate of care-giving, lower population turnover, and lower increases in average population intelligence. Overall, these simulations suggest that diseases that are most likely to have led to the evolution of care-giving in the human lineage were those with low costs to caregivers which persisted at a prevalence low enough not to disrupt the kin networks along which care was provided. Although only healthy agents could give care and reproduce in our model, high rates of costly care-giving may not be expected if kin have sublethal diseases that do not reduce their reproductive success.

It is noteworthy that for all three diseases that produced care-giving, the final rate of care-giving was low, with a maximum of 6.7% of the diseased receiving care under measles-like conditions. Furthermore, a recovery rate of only 50% after care suggests that over the course of hominin evolution even low rates of relatively ineffective care may have been sufficient to select for increasing intelligence and disease recognition. We expect that the first forms of care-giving among hominins would have included assistance with hygiene, such as cleaning of wounds and topical infections (Dittus and Ratnayeke 1989) and provisioning with food and water (Sugiyama 2004). These mechanisms would not have required an understanding of disease processes and could have piggybacked on basal social grooming behaviors observed in nonhuman primates (Dittus and Ratnayeke 1989) and communal provisioning behaviors that may have evolved during the evolution of cooperative breeding (Burkart et al. 2009; Gurven et al. 2000; Hawkes 2003; Hill et al. 2009; Hrdy 2009; Sugiyama 2004; Sugiyama and Chacon 2000).

## *Model 2*

The Model 2 results demonstrate that avoidance alone does not select for greater cognitive abilities. Avoidance produced no net change in average population intelligence in

the scabies-like and measles-like conditions and a *decrease* in average population intelligence in for the CCHF-like and Ebola-like diseases. The scabies-like and measles-like diseases produced higher population sizes and disease prevalences *above* 50%, thus an agent who moves away from infected kin is likely to encounter other infected individuals. This results in a lack of selection for disease recognition and avoidance. In contrast, the CCHF-like and Ebola-like diseases produced lower population sizes and prevalences *below* 50%, thus an agent who avoids infected kin is less likely to encounter other infected agents. This results in selection to isolate oneself. The most efficient way for agents to isolate themselves in a population with a prevalence under 50%, is to miscategorize healthy individuals as ill, thus triggering avoidance. Because lower intelligence agents have less accurate disease recognition, this produces selection to *decrease* intelligence.

These findings are relevant for species that do not give care. It suggests that avoidance of high prevalence, low fatality diseases is likely to be an ineffective strategy. As a result these diseases do not exert selection for or against cognitive abilities under an avoidance only paradigm. In contrast, avoidance is an effective strategy against low prevalence, high fatality diseases producing selection for avoidance behavior and selection against sociality.

### *Implications of care-giving*

A comparison of the results from Model 1 (care-giving model) with Model 2 (avoidance model) indicates that care-giving alters the progression of the disease through the population. For the scabies-like and measles-like diseases, care-giving resulted in significantly higher population sizes and lower prevalences than an avoidance only strategy. Thus for these diseases, which are the two diseases for which care was given to both close and distant kin ( $r=0.5$  and  $r=0.25$ , respectively), care-giving served to control the disease in the population.

Two of the diseases, the measles-like and the CCHF-like diseases, show distinct cycles of disease outbreaks and population crashes (“boom and bust” dynamic, Fig. 2-3). The lack of congruence between the relatively constant slope of the intelligence curves (Fig. 4) and the boom-bust oscillations of population size and prevalence, is a reflection of the fact that selection on intelligence is occurring throughout the boom-bust cycle and not intermittently only when specific conditions are met (e.g., a particular population size or prevalence). This dynamic is quantified through the interaction term of the mixed model analysis in which intelligence increases are the result of complex interactions between prevalence and population size. Because the two diseases progress differently through the population, they also exert selection on intelligence in slightly different ways. The measles-like disease produces one oscillation of the boom-bust outbreak cycle of population and prevalence peaks and crashes; the CCHF-like disease produces multiple, more rapid oscillations.

The measles-like disease shows a very pronounced “bust” phase early in the run. Population size is high when the disease is first introduced (Fig. 2B, Model 1 curve). This produces a high rate of care-giving and strong selection for intelligence (left panel, Fig. 6B). As the prevalence increases (Fig. 3B, Model 1 curve), low intelligence matrilineal lines recognize diseased kin less accurately, and provide less successful care, causing them to succumb to the disease. This produces a decrease in population size and an increase in average population intelligence (Fig. 4B, Model 1 curve). At high prevalences, selection for intelligence is maintained regardless of the population size (right panel, Fig. 6B). Intelligence plateaus about half way through the run when the population size rebounds slightly but remains low and prevalence decreases slightly from its earlier peak and remains moderate. With a low population size, intermediate prevalence, and a decreased rate of care-giving (Fig. 1B, measles-like curve), the population maintains the higher intelligence, but does not continue to

increase it (change in intelligence approaches 0 in left side of middle panel, Fig. 6B).

Intelligence plateaus as the boom-bust outbreak oscillations cease.

The CCHF-like disease produces a very pronounced boom-bust cycle with several peaks and crashes in population size and prevalence. Selection for increasing intelligence occurs both during low population sizes and low prevalences (left panel, Fig. 7B) and during high population size and high prevalences (right panel, Fig. 7B). When the boom-bust dynamic stops about halfway through the run and the population stabilizes at intermediate population sizes and prevalences, intelligence plateaus (Figs. 2C, 3C, 4C Model 1 curves and middle panel, Fig. 7B).

Interestingly, when the population infected with the measles-like disease engages in care-giving, it experiences less pronounced oscillations of the “boom and bust” outbreak cycle (Fig. 3) indicating that care-giving serves to control the spread of the disease through the population. Because of the higher risks of providing care under the CCHF-like conditions, only close kin ( $r=0.5$ ) receive care. This lower level of care is less effective at controlling the spread of the disease, perhaps suggesting that a certain threshold must be achieved in order to disrupt the boom-bust outbreak cycle (boom-bust dynamics: (Keeling and Grenfell 1997)). Alternatively, the higher fatality rate and more rapid transmission of the CCHF-like diseases produces faster outbreak cycles, which may make it more difficult for care-giving to disrupt the boom-bust outbreak cycle even though it still selects for increasing cognitive abilities.

For both the measles-like and CCHF-like diseases, the most pronounced outbreaks occur early in the model run, which is also when the greatest increases in intelligence are occurring (Fig. 6A and 7A). In the second half of the run, when the boom-bust dynamic is less pronounced, intelligence plateaus. This suggests that over the course of human evolution, sustained increases in intelligence may have occurred through repeated introductions of novel diseases into naïve populations. The greatest selection would have occurred shortly after the



introduction when the disease was spreading and care-giving behavior had not yet managed to reduce the size of the outbreaks and subsequent population crashes.

### *Significance for human evolution*

Our model was parameterized based upon group sizes, spatial scales, and population densities derived from the fossil record and modern foraging peoples (Grove et al. 2012; Layton et al. 2012). Our goal was not to recreate a particular hominin population, but to explore the effects of different disease characteristics on the evolution of care-giving and increased cognition in a population with hominin characteristics.

We created an SIS model (susceptible-infected-susceptible) where recovered individuals are just as susceptible as those who were never infected. However, for many diseases, recovered individuals are temporarily or permanently immune to re-infection, potentially increasing their ability to provide care. We expect that immunity would increase the rate of care-giving. Diseases likely to select for care-giving among kin may be diseases which frequently infect children and then convey lifetime immunity. Under this scenario, adults who survived to reproduce would have extensive knowledge of the disease's symptoms, making recognition likely, and the immunity to enable them to provide effective care. Several well-known childhood diseases that follow this pattern (e.g., measles, smallpox) have been dated to the origins of agriculture, animal domestication, and the subsequent population increases (Harper and Armelagos 2013). However, as more genetic studies are conducted, increasing numbers of pathogens are showing pre-agricultural origins, including some that were previously believed to be post-agricultural (e.g., tapeworms, TB (Harper and Armelagos 2013; Hoberg et al. 2001; Hurtado et al. 2008; Stone et al. 2009). Tapeworms, TB, typhoid fever, whooping cough, and Epstein Barr virus, among others, have been shown to predate agriculture (Harper and Armelagos 2013; Hoberg et al. 2001; Hurtado et al. 2008; Stone et al.

2009), suggesting that ancestral hominins harbored significant numbers of infectious diseases. Based on our models, diseases with low risks to care-givers, high inclusive fitness pay-offs for care-givers, and prevalences low enough not to disrupt the kin networks along which care could be given would have exerted the strongest selection for increased cognition. Through repeated introductions of novel diseases over millions of years, such diseases could have selected for accurate disease recognition, increased care-giving among kin, and produced the social and cognitive origins of human medical care.

### **A novel hypothesis of human cognitive evolution and future directions**

Our novel hypothesis of primate, including human cognitive evolution, is *not* mutually exclusive with the social brain hypothesis (Dunbar 1998). As social species evolved the cognitive capacities for social cognition, such as processing information gleaned from faces (Leopold and Rhodes 2010; Sartori et al. 2011), voices (Belin 2006; Belin et al. 2004), and movement patterns (Loula et al. 2005; Peterman et al. 2014; Sartori et al. 2011), they may have also obtained the ability to use this information to recognize disease symptoms. They could detect changes in facial coloration and texture due to fever or rashes, changes in vocalizations due to coughing, nasal discharge or reduced lung capacity, and changes in movement/gait due to weakness, lethargy, or signs of pain (Chapman et al. 2005; Fink and Matts 2008; Hart 1988). The proximate mechanisms are relatively simple in that they do *not* require individuals to have an abstract concept of “disease.” Instead, individuals that are able to accurately recognize disease would have increased fitness due to being able to avoid infectious individuals or provide care to kin. Though studies of disease recognition in nonhuman animals are relatively rare, several species do appear to recognize the health status of conspecifics, i.e., social lobsters (Behringer et al. 2006), pipefish (Rosenqvist and

Johansson 1995), bullfrog tadpoles (Kiesecker et al. 1999), rodents (Kavaliers et al. 1997), house finches (Bouwman and Hawley 2010; Zylberberg et al. 2012), but see (Nunn 2003).

We predict that as hominin social complexity increased, i.e., group sizes, social network sizes, frequencies of cooperation and social learning, etc. (Aiello and Dunbar 1993; Burkart et al. 2014; Burkart et al. 2009; Dunbar 1998; Gamble et al. 2011; Grove et al. 2012; Layton et al. 2012; Tomasello 2014), hominins would have substantially increased their risk of disease transmission, producing heightened selection for disease recognition and care-giving. We make several predictions that enable paleoanthropologists, archaeologists, primatologists, human ecologists, geneticists and immunologists to test our novel hypothesis of human cognitive evolution:

- 1) Humans and nonhuman primates have very similar disease profiles in that we share many of the same diseases with viral, bacterial, and gastrointestinal parasitic zoonoses occurring from nonhuman primates to humans and vice versa (Chapman et al. 2005; Jones et al. 2008; Lloyd-Smith et al. 2009; Wolfe et al. 2007). However, what has received very little attention is how humans and nonhuman primates may differ in the expression of disease symptoms. Humans, relative to nonhuman primates have much less body hair. Though our nakedness may reduce ectoparasite load (Pagel and Bodmer 2003; Weiss 2007), it also provides a visually unobstructed surface for displaying rashes, lesions, swelling, and inflammation, and bruising. Humans, relative to nonhuman primates, also have white scleras around their eyes, a signal that has been argued to draw attention to gaze direction (Kobayashi and Kohshima 2001; Tomasello et al. 2007), but also turns a dramatic “bloodshot” red when we are under emotional stress or ill (Provine et al. 2011). **Prediction 1:** *If humans have been selected to solicit care from others, they should display exaggerated signals of ill*

health, relative to nonhuman primates experiencing the same disease and degree of morbidity/mortality.

- 2) It is becoming increasingly possible to date the origins of many diseases afflicting humans i.e., (Harper and Armelagos 2013; Stone et al. 2009). As more accurate dates are obtained for more diseases, it will be possible to examine whether hominin populations carried an increased disease load as they increased in social complexity. Social complexity could be operationalized in the fossil record through the brain size – group size relationship (Aiello and Dunbar 1993; Dunbar 1998; Gamble et al. 2011; Grove et al. 2012; Layton et al. 2012), through evidence of increased behavioral and technological complexity in the archaeological record (Gowlett et al. 2012; Shultz et al. 2012), or through fossil evidence for the shift to cooperative breeding (Aiello and Key 2002; Shultz et al. 2012). **Prediction 1:** *If larger hominin communities sustained greater disease loads, then periods of rapidly increasing community sizes (operationalized with expanding brain sizes (Aiello and Dunbar 1993; Dunbar 1998; Gamble et al. 2011; Grove et al. 2012)) should coincide with the evolution of diseases new to hominins.* **Prediction 2:** *If social learning/cooperation lead to increased disease transmission (McCabe et al. 2015), then increasing behavioral/technological complexity in the archaeological record (Gamble et al. 2011; Gowlett et al. 2012; Shultz et al. 2012) should coincide with the evolution of diseases new to hominins.* **Prediction 3:** *If cooperatively breeding increased disease transmission, then evidence for cooperative breeding in the fossil record (Aiello and Key 2002; Shultz et al. 2012) should coincide with the evolution of diseases new to hominins, particularly those that afflict children.* These predictions are not mutually exclusive. According to the results of our model, we would expect a high proportion of these diseases to present low costs and high fitness payoffs to care-givers and persist at prevalences that are low

enough not to disrupt the kin networks along which care is provided. Possibilities include infections that leave survivors immune.

- 3) An additional avenue for examining the role of disease during the evolution of human social complexity would be through cross-species comparisons of immune investment. If hominins have experienced an unusually high rate of disease exposure, either through their extensive social networks or through providing care to diseased kin, they may have invested heavily in immune defenses. Recent work on introgression between anatomically modern humans (AMH) and neandertals has proposed that one of the major advantages may have been the acquisition of novel immune genes from neandertals as AMH expanded northward into novel environments and encountered novel pathogens (Houldcroft and Underdown 2016). Prior studies indicate that there are cross-species differences in immune investment according to mating system (but not group size or density in primates) (Nunn et al. 2000), the risk of environmentally transmitted parasites and injuries due to predator attacks in anthropoids (Semple et al. 2002), coloniality in birds (Moller et al. 2001), and cooperative breeding in birds (Spottiswoode 2008). **Prediction 1:** *If hominins' increased social complexity required them to invest heavily in immune defenses, the human immune system should show similar adaptations to other species that have extremely large social networks and high interaction rates. Prediction 2:* *If the evolution of cooperative breeding required hominins to invest heavily in immune defenses, then the human immune system should show similar adaptations to other cooperatively breeding species. Prediction 3:* *If the evolution of providing care to diseased conspecifics required hominins to invest heavily in immune defenses, the human immune system should show adaptations that are either extreme or unusual.* (These predictions are not mutually exclusive). While many of the earlier studies were

done with white blood cell counts, i.e., (Nunn et al. 2000), the field of ecological immunology is growing rapidly with new techniques being continually developed (Downs et al. 2014; Larsen et al. 2014). This should make it increasingly possible to parse out how different selective forces may have acted on different elements of a species' immune system.

## **Conclusions**

Our model indicates that disease circulating amongst kin groups can select for care-giving among kin and greater cognitive abilities. Moreover, the characteristics of the diseases can generate different strengths of selection. Diseases with lower costs and higher pay offs produced stronger selection, yielding higher care-giving rates and greater increases in average population intelligence.

When a care-giving strategy was compared with an avoidance only strategy, the care-giving strategy controlled the transmission of the disease through the population by reducing the severity of disease outbreaks and population crashes. Because this cycle of outbreaks and population crashes was associated with the most rapid increases in intelligence, we propose that the repeated introduction of novel diseases into naïve populations may have led to sustained selection for increasing disease recognition and cognitive abilities throughout human evolution. Moreover, the unique ability of hominins to control the spread of disease through care-giving behaviors may have facilitated increased social complexity, and ultimately lead to the evolution of medical care in humans. Finally, we set out predictions derived from our disease recognition hypothesis of hominin cognitive evolution that can be tested by paleoanthropologists, archaeologists, geneticists, and primatologists.

## **Data accessibility**

The ODD descriptions of Model 1 (caregiving) and Model 2 (avoidance only) are found in Appendices A and B, respectively. The code is available in supplementary files 1 and 2, respectively. The files containing the code can be opened with standard text editing programs such as WordPad.

### **Conflict of interests**

None.

### **Authors' contributions**

SEK designed the study, programmed the model, analyzed the data, and wrote the manuscript. TRB and CAC contributed to all stages. RWB contributed to the development of the ideas and manuscript preparation.

### **Acknowledgements**

This work was funded by the Fonds de Recherche du Québec – Nature et Technologies and the Québec Center for Biodiversity Science. The funders were not involved in the study design, the collection, analysis and interpretation of data, writing of the report or decision to submit it.

### **References**

- Aiello LC, Dunbar R (1993) Neocortex size, group size, and the evolution of language. *Curr Anthropol* 34:184-193
- Aiello LC, Key C (2002) Energetic consequences of being a *Homo erectus* Female. *Am J Hum Biol* 14:551-565
- Alizon S, Hurford A, Mideo N, Van Baalen M (2009) Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *J Evol Biol* 22:245-259
- Anderson JR, Gillies A, Lock LC (2010) Pan thanatology. *Curr Biol* 20:R349-R351

- 961 Armelagos GJ, Brown PJ, Turner B (2005) Evolutionary, historical and political economic perspectives  
962 on health and disease. *Soc Sci Med* 61:755-765
- 963 Atkinson QD, Gray RD, Drummond A (2008) mtDNA variation predicts population size in humans and  
964 reveals a major southern Asian chapter in human prehistory. *Mol Biol Evol* 25:468-474
- 965 Aylward B, Barboza P, Bawo L, Bertherat E, Bilivogui P, Blake I, Brennan R, Briand S, Chakauya JM,  
966 Chitala K, Conteh RM, Cori A, Croisier A, Dangou JM, Diallo B, Donnelly CA, Dye C, Eckmanns T,  
967 Ferguson NM, Formenty P, Fuhrer C, Fukuda K, Garske T, Gasasira A, Gbanyan S, Graaff P, Heleze E,  
968 Jambai A, Jombart T, Kasolo F, Kadiobo AM, Keita S, Kertesz D, Kone M, Lane C, Markoff J, Massaquoi  
969 M, Mills H, Mulba JM, Musa E, Myhre J, Nasidi A, Nilles E, Nouvellet P, Nshimirimana D, Nuttall I,  
970 Nyenswah T, Olu O, Pendergast S, Perea W, Polonsky J, Riley S, Ronveaux O, Sakoba K, Krishnan RSG,  
971 Senga M, Shuaib F, Van Kerkhove MD, Vaz R, Kannangarage NW, Yoti Z (2014) Ebola virus disease in  
972 West Africa - The first 9 months of the epidemic and forward projections. *N Engl J Med* 371:1481-  
973 1495
- 974 Beamish EK, O'Riain MJ (2014) The effects of permanent injury on the behavior and diet of  
975 commensal chacma baboons (*Papio ursinus*) in the Cape Peninsula, South Africa. *Int J Primatol*  
976 35:1004-1020
- 977 Behringer DC, Butler MJ, Shields JD (2006) Avoidance of disease by social lobsters. *Nature* 441:421-  
978 421
- 979 Belin P (2006) Voice processing in human and non-human primates. *Philos Trans R Soc Lond B Biol*  
980 *Sci* 361:2091-2107
- 981 Belin P, Fecteau S, Bedard C (2004) Thinking the voice: neural correlates of voice perception. *Trends*  
982 *in Cognitive Sciences* 8:129-135
- 983 Bezerra BM, Keasey MP, Schiel N, Souto AD (2014) Responses towards a dying adult group member  
984 in a wild New World monkey. *Primates* 55:185-188
- 985 Bonmati A, Gomez-Olivencia A, Arsuaga JL, Carretero JM, Gracia A, Martinez I, Lorenzo C, de Castro  
986 JMB, Carbonell E (2010) Middle Pleistocene lower back and pelvis from an aged human individual  
987 from the Sima de los Huesos site, Spain. *Proc Natl Acad Sci U S A* 107:18386-18391
- 988 Bouwman KM, Hawley DM (2010) Sickness behaviour acting as an evolutionary trap? Male house  
989 finches preferentially feed near diseased conspecifics. *Biology Letters* 6:462-465
- 990 Burkart J, Allon O, Amici F, Fichtel C, Finkenwirth C, Heschl A, Huber L, Isler K, Kosonen ZK, Martins E,  
991 Meulman EJ, Richiger R, Rueth K, Spillmann B, Wiesendanger S, van Schaik C (2014) The evolutionary  
992 origin of human hyper-cooperation. *Nature Communications* 5:4747
- 993 Burkart JM, Hrdy SB, Van Schaik CP (2009) Cooperative breeding and human cognitive evolution.  
994 *Evol Anthropol* 18:175-186
- 995 Byrne RW, Bates LA (2007) Sociality, evolution and cognition. *Curr Biol* 17:R714-R723
- 996 Byrne RW, Stokes EJ (2002) Effects of manual disability on feeding skills in gorillas and chimpanzees.  
997 *Int J Primatol* 23:539-554
- 998 Chapais B, Berman CM (eds) (2004) *Kinship and Behavior in Primates*. Oxford University Press, New  
999 York



- 1000 Chapais B, Gauthier C, PrudHomme J, Vasey P (1997) Relatedness threshold for nepotism in  
1001 Japanese macaques. *Anim Behav* 53:1089-1101
- 1002 Chapman C, Gillespie TR, Goldberg TL (2005) Primates and the ecology of their infectious diseases:  
1003 How will anthropogenic change affect host-parasite interactions? *Evol Anthropol* 14:134-144
- 1004 Coqueugniot H, Dutour O, Arensburg B, Duda H, Vandermeersch B, Tillier AM (2014) Earliest cranio-  
1005 encephalic trauma from the Levantine Middle Palaeolithic: 3D reappraisal of the Qafzeh 11 skull,  
1006 consequences of pediatric brain damage on individual life condition and social care. *Plos One* 9:10
- 1007 Crubezy E, Trinkaus E (1992) Shanidar 1: A case of hyperostotic disease (DISH) in the Middle  
1008 Paleolithic. *Am J Phys Anthropol* 89:411-420
- 1009 Cuozzo FP, Sauter ML (2004) Tooth loss, survival, and resource use in wild ring-tailed lemurs (*Lemur*  
1010 *catta*): implications for inferring conspecific care in fossil hominids. *J Hum Evol* 46:623-631
- 1011 Davenport LC (2010) Aid to a declining matriarch in the giant otter (*Pteronura brasiliensis*). *Plos One*  
1012 5:6
- 1013 Deaner RO, van Schaik C, Johnson V (2006) Do some taxa have better domain-general cognition than  
1014 others? A meta-analysis of nonhuman primate studies. *Evolutionary Psychology* 4:149-196
- 1015 DeGusta D (2002) Comparative skeletal pathology and the case for conspecific care in Middle  
1016 Pleistocene hominids. *J Archaeol Sci* 29:1435-1438
- 1017 DeGusta D (2003) Aubesier 11 is not evidence of Neanderthal conspecific care. *J Hum Evol* 45:91-94
- 1018 Dettwyler KA (1991) Can Paleoanthropology provide evidence for compassion? *Am J Phys Anthropol*  
1019 84:375-384
- 1020 Dittus WPJ, Ratnayeke SM (1989) Individual and social behavioral-responses to injury in wild Toque  
1021 macaques (*Macaca sinica*). *Int J Primatol* 10:215-234
- 1022 Douglas-Hamilton I, Bhalla S, Wittemyer G, Vollrath F (2006) Behavioural reactions of elephants  
1023 towards a dying and deceased matriarch. *Appl Anim Behav Sci* 100:87-102
- 1024 Downs CJ, Adelman JS, Demas GE (2014) Mechanisms and methods in ecoimmunology: Integrating  
1025 within-organism and between-organism processes. *Integrative and Comparative Biology* 54:340-352
- 1026 Dunbar R (1998) The social brain hypothesis. *Evol Anthropol* 6:178-190
- 1027 Ewald PW (1993) The evolution of virulence. *Sci Am* 268:86-93
- 1028 Fernandes HBF, Woodley MA, te Nijenhuis J (2014) Differences in cognitive abilities among primates  
1029 are concentrated on G: Phenotypic and phylogenetic comparisons with two meta-analytical  
1030 databases. *Intelligence* 46:311-322
- 1031 Field A (2013) *Discovering Statistics Using IBM SPSS Statistics*, 4th edn. Sage, London
- 1032 Fink B, Mates PJ (2008) The effects of skin colour distribution and topography cues on the perception  
1033 of female facial age and health. *J Eur Acad Dermatol Venereol* 22:493-498

- 1034 Gamble C, Gowlett J, Dunbar R (2011) The social brain and the shape of the palaeolithic. Cambridge  
1035 Archaeological Journal 21:115-135
- 1036 Gowlett J, Gamble C, Dunbar R (2012) Human Evolution and the Archaeology of the Social Brain. Curr  
1037 Anthropol 53:693-722
- 1038 Gracia A, Arsuaga JL, Martínez I, Lorenzo C, Carretero JM, Bermúdez de Castro JM, Carbonell E  
1039 (2009) Craniosynostosis in the middle pleistocene human cranium 14 from the Sima de los Huesos,  
1040 Atapuerca, Spain. Proceedings of the National Academy of Sciences 106:6573-6578
- 1041 Griffin R, Nunn C (2012) Community structure and the spread of infectious disease in primate social  
1042 networks. Evol Ecol 26:779-800
- 1043 Grove M, Pearce E, Dunbar RIM (2012) Fission-fusion and the evolution of hominin social systems. J  
1044 Hum Evol 62:191-200
- 1045 Guo DM, Li KC, Peters TR, Snively BM, Poehling KA, Zhou XB (2015) Multi-scale modeling for the  
1046 transmission of influenza and the evaluation of interventions toward it. Scientific Reports 5:9
- 1047 Gurven M, Allen-Arave W, Hill K, Hurtado M (2000) "It's a Wonderful Life": signaling generosity  
1048 among the Ache of Paraguay. Evolution and Human Behavior 21:263-282
- 1049 Haeusler M, Schiess R, Boeni T (2013) Evidence for juvenile disc herniation in a *Homo erectus* boy  
1050 skeleton. Spine 38:E123-E128
- 1051 Hamilton WD (1964) The genetical evolution of social behavior. I and II. J Theor Biol 7:1-52
- 1052 Harper KN, Armelagos GJ (2013) Genomics, the origins of agriculture, and our changing microbe-  
1053 scape: Time to revisit some old tales and tell some new ones. Am J Phys Anthropol 152:135-152
- 1054 Hart BL (1988) Biological basis of the behavior of sick animals. Neurosci Biobehav Rev 12:123-137
- 1055 Hart BL (2011) Behavioural defences in animals against pathogens and parasites: parallels with the  
1056 pillars of medicine in humans. Philos Trans R Soc Lond B Biol Sci 366:3406-3417
- 1057 Hatchwell BJ (2010) Cryptic kin selection: Kin structure in vertebrate populations and opportunities  
1058 for kin-directed cooperation. Ethology 116:203-216
- 1059 Hawkes K (2003) Grandmothers and the evolution of human longevity. Am J Hum Biol 15:380-400
- 1060 Herrmann E, Call J, Hernández-Lloreda MV, Hare B, Tomasello M (2007) Humans have evolved  
1061 specialized skills of social cognition: The cultural intelligence hypothesis. Science 317:1360
- 1062 Hill K, Barton M, Hurtado A (2009) The emergence of human uniqueness: Characters underlying  
1063 behavioral modernity. Evol Anthropol 18:187-200
- 1064 Hill KR, Walker RS, Bozicevic M, Eder J, Headland T, Hewlett B, Hurtado AM, Marlowe FW, Wiessner  
1065 P, Wood B (2011) Co-residence patterns in hunter-gatherer societies show unique human social  
1066 structure. Science 331:1286-1289
- 1067 Hill KR, Wood BM, Baggio J, Hurtado AM, Boyd RT (2014) Hunter-gatherer inter-band interaction  
1068 rates: Implications for cumulative culture. Plos One 9:9

- 1069 Hoberg EP, Alkire NL, de Queiroz A, Jones A (2001) Out of Africa: Origins of the *Taenia* tapeworms in  
1070 humans. P Roy Soc B-Biol Sci 268:781-787
- 1071 Houldcroft CJ, Underdown SJ (2016) Neanderthal genomics suggests a pleistocene time frame for  
1072 the first epidemiologic transition. Am J Phys Anthropol 160:379-388
- 1073 Hrdy SB (2009) Mothers and Others: The Evolutionary Origins of Mutual Understanding. Belknap  
1074 Press of Harvard University Press, Cambridge
- 1075 Hublin JJ (2009) The prehistory of compassion. Proceedings of the National Academy of Sciences  
1076 106:6429-6430
- 1077 Hurtado AM, Frey MA, Hurtado I, Hill K, Baker J (2008) The role of helminthes in human evolution  
1078 implications for global health in the 21st century. Society for the Study of Human Biology Series  
1079 48:153-180
- 1080 Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P (2008) Global trends in  
1081 emerging infectious diseases. Nature 451:990-994
- 1082 Kahm M, Hasenbrink G, Lichtenberg-Fraté H, Ludwig J, Kschischo M (2010) grofit: Fitting biological  
1083 growth curves with R. Journal of Statistical Software 33:1-21
- 1084 Kavaliers M, Colwell D, Ossenkopp KP, Perrot-Sinal TS (1997) Altered responses to female odors in  
1085 parasitized male mice: Neuromodulatory mechanisms and relations to female choice. Behav Ecol  
1086 Sociobiol 40:373-384
- 1087 Keeling MJ, Grenfell BT (1997) Disease extinction and community size: Modeling the persistence of  
1088 measles. Science 275:65-67
- 1089 Kiesecker JM, Skelly DK, Beard KH, Preisser E (1999) Behavioral reduction of infection risk. Proc Natl  
1090 Acad Sci U S A 96:9165-9168
- 1091 Kobayashi H, Kohshima S (2001) Unique morphology of the human eye and its adaptive meaning:  
1092 comparative studies on external morphology of the primate eye. J Hum Evol 40:419-435
- 1093 Larsen PA, Campbell CR, Yoder AD (2014) Next-generation approaches to advancing eco-  
1094 immunogenomic research in critically endangered primates. Molecular Ecology Resources 14:1198-  
1095 1209
- 1096 Layton R, O'Hara S, Bilsborough A (2012) Antiquity and social functions of multilevel social  
1097 organization among human hunter-gatherers. Int J Primatol 33:1215-1245
- 1098 Lebel S, Trinkaus E (2002) Middle Pleistocene human remains from the Bau de l'Aubiesier. J Hum Evol  
1099 43:659-685
- 1100 Lebel S, Trinkaus E, Faure M, Fernandez P, Guérin C, Richter D, Mercier N, Valladas H, Wagner GA  
1101 (2001) Comparative morphology and paleobiology of Middle Pleistocene human remains from the  
1102 Bau de l'Aubiesier, Vaucluse, France. Proceedings of the National Academy of Sciences 98:11097-  
1103 11102
- 1104 Lee JJ (2007) A g beyond Homo sapiens? Some hints and suggestions. Intelligence 35:253-265
- 1105 Leopold DA, Rhodes G (2010) A comparative view of face perception. J Comp Psychol 124:233-251

- 1106 Lloyd-Smith JO, George D, Pepin KM, Pitzer VE, Pulliam JRC, Dobson AP, Hudson PJ, Grenfell BT  
1107 (2009) Epidemic Dynamics at the Human-Animal Interface. *Science* 326:1362-1367
- 1108 Lordkipanidze D, Vekua A, Ferring R, Rightmire GP, Agustill J, Kiladze G, Mouskhelishvili A, Nioradze  
1109 M, de Leon MSP, Tappen M, Zollikofer CPE (2005) The earliest toothless hominin skull. *Nature*  
1110 434:717-718
- 1111 Lordkipanidze D, Vekua A, Ferring R, Rightmire GP, Zollikofer CPE, De Leon MSP, Agusti J, Kiladze G,  
1112 Mouskhelishvili A, Nioradze M, Tappen M (2006) A fourth hominin skull from Dmanisi, Georgia.  
1113 *Anatomical Record Part a-Discoveries in Molecular Cellular and Evolutionary Biology* 288A:1146-  
1114 1157
- 1115 Loula F, Prasad S, Harber K, Shiffrar M (2005) Recognizing people from their movement. *Journal of*  
1116 *Experimental Psychology-Human Perception and Performance* 31:210-220
- 1117 McBrearty S, Brooks AS (2000) The revolution that wasn't: A new interpretation of the origin of  
1118 modern human behavior. *J Hum Evol* 39:453-563
- 1119 McCabe CM, Reader SM, Nunn CL (2015) Infectious disease, behavioural flexibility and the evolution  
1120 of culture in primates. *P Roy Soc B-Biol Sci* 282:9
- 1121 Merler S, Ajelli M, Fumanelli L, Gomes MFC, Piontti APY, Rossi L, Chao DL, Longini IM, Halloran ME,  
1122 Vespignani A (2015) Spatiotemporal spread of the 2014 outbreak of Ebola virus disease in Liberia  
1123 and the effectiveness of non-pharmaceutical interventions: a computational modelling analysis.  
1124 *Lancet Infectious Diseases* 15:204-211
- 1125 Moller AP, Merino S, Brown CR, Robertson RJ (2001) Immune Defense and Host Sociality: A  
1126 Comparative Study of Swallows and Martins. *Am Nat* 158:136-145
- 1127 Montgomery PQ, Williams HOL, Reading N, Stringer CB (1994) An assessment of the temporal bone  
1128 lesions of the Broken Hill cranium. *J Archaeol Sci* 21:331-337
- 1129 Munn J (2006) Effects of injury on the locomotion of free-living chimpanzees in the Bodongo Forest  
1130 Reserve, Uganda. In: Newton-Fisher NE, Notman H, Paterson JD, Reynolds V (eds) *Primates of*  
1131 *Western Uganda*. Springer Science+Business Media, New York, pp 259-280
- 1132 Nakamichi M, Koyama N, Jolly A (1996) Maternal responses to dead and dying infants in wild troops  
1133 of ring-tailed lemurs at the Berenty reserve, Madagascar. *Int J Primatol* 17:505-523
- 1134 Neisser U (1967) *Cognitive Psychology*, 1st edn. Prentice Hall, Englewood Cliffs
- 1135 Nunn C, Altizer S (2006) *Infectious Diseases in Primates: Behavior, Ecology and Evolution*. Oxford  
1136 University Press, New York
- 1137 Nunn CL (2003) Behavioural defenses against sexually transmitted diseases in primates. *Anim Behav*  
1138 66:37-48
- 1139 Nunn CL, Gittleman JL, Antonovics J (2000) Promiscuity and the primate immune system. *Science*  
1140 290:1168-1170
- 1141 Pagel M, Bodmer W (2003) A naked ape would have fewer parasites. *P Roy Soc B-Biol Sci* 270:S117-  
1142 S119

- 1143 Park KJ, Sohn H, An YR, Moon DY, Choi SG, An DH (2013) An unusual case of care-giving behavior in  
1144 wild long-beaked common dolphins (*Delphinus capensis*) in the East Sea. *Mar Mamm Sci* 29:E508-  
1145 E514
- 1146 Peterman JS, Christensen A, Giese MA, Park S (2014) Extraction of social information from gait in  
1147 schizophrenia. *Psychol Med* 44:987-996
- 1148 Powell A, Shennan S, Thomas MG (2009) Late Pleistocene demography and the appearance of  
1149 modern human behavior. *Science* 324:1298-1301
- 1150 Provine RR, Cabrera MO, Brocato NW, Krosnowski KA (2011) When the Whites of the Eyes are Red: A  
1151 Uniquely Human Cue. *Ethology* 117:395-399
- 1152 Railsback SF, Grimm V (2011) Agent-Based and Individual-Based Modeling: A Practical Introduction.  
1153 Princeton University Press, Princeton
- 1154 RCoreTeam (2011) R: A language and environment for statistical computing R Foundation for  
1155 Statistical Computing, Viena, Austria
- 1156 RCoreTeam (2015) R: A language and environment for statistical computing. R Foudnation for  
1157 Statistical Computing, Vienna, Austria
- 1158 Reader SM, Hager Y, Laland KN (2011) The evolution of primate general and cultural intelligence.  
1159 *Philosophical Transactions of the Royal Society B: Biological Sciences* 366:1017-1027
- 1160 Rendall D (2004) Recognizing kin: mechanisms, media, minds, modules, and muddles. In: Chapais B,  
1161 Berman CM (eds) *Kinship and Behavior in Primates*. Oxford University Press, New York, pp 295-316
- 1162 Rifkin JL, Nunn CL, Garamszegi LZ (2012) Do animals living in larger groups experience greater  
1163 parasitism? A meta-analysis. *Am Nat* 180:70-82
- 1164 Rosenqvist G, Johansson K (1995) Male avoidance of parasitized females explained by direct benefits  
1165 in a pipefish. *Anim Behav* 49:1039-1045
- 1166 RStudio (2014) RStudio: Integrated development environment for R Boston, MA
- 1167 Sartori L, Becchio C, Castiello U (2011) Cues to intention: The role of movement information.  
1168 *Cognition* 119:242-252
- 1169 Schiess R, Boeni T, Rühli F, Haeusler M (2014) Revisiting scoliosis in the KNM-WT 15000 *Homo*  
1170 *erectus* skeleton. *J Hum Evol* 67:48-59
- 1171 Semple S, Cowlshaw G, Bennett PM (2002) Immune system evolution among anthropoid primates:  
1172 parasites, injuries and predators. *P Roy Soc B-Biol Sci* 269:1031-1037
- 1173 Shang H, Trinkaus E (2008) An ectocranial lesion on the Middle Pleistocene human cranium from  
1174 Hulu Cave, Nanjing, China. *Am J Phys Anthropol* 135:431-437
- 1175 Sherman PW (1988) Levels of analysis. *Anim Behav* 36:616-619
- 1176 Shultz S, Nelson E, Dunbar RIM (2012) Hominin cognitive evolution: identifying patterns and  
1177 processes in the fossil and archaeological record. *Philos Trans R Soc Lond B Biol Sci* 367:2130-2140

- 1178 Silk JB (2009) Nepotistic cooperation in non-human primate groups. *Philos Trans R Soc Lond B Biol*  
1179 *Sci* 364:3243-3254
- 1180 Spottiswoode CN (2008) Cooperative breeding and immunity: a comparative study of PHA response  
1181 in African birds. *Behav Ecol Sociobiol* 62:963-974
- 1182 Stokes EJ, Byrne R (2006) Effect of snare injuries on the fig-feeding behavior of chimpanzees of the  
1183 Bondogo Forest, Uganda: Behavioral adaptations and long-term implications. In: Newton-Fisher NE,  
1184 Notman H, Paterson JD, Reynolds V (eds) *Primates of Western Uganda*. Springer Science+Business  
1185 Media, New York, pp 281-297
- 1186 Stone AC, Wilbur AK, Buikstra JE, Roberts CA (2009) Tuberculosis and leprosy in perspective. *Yearb*  
1187 *Phys Anthropol* 52:66-94
- 1188 Sugiyama LS (2004) Illness, injury, and disability among Shiwiari forager-horticulturists: Implications  
1189 of health-risk buffering for the evolution of human life history. *Am J Phys Anthropol* 123:371-389
- 1190 Sugiyama LS, Chacon R (2000) Effects of illness and injury on foraging among the Yora and Shiwiari:  
1191 Pathology risk as adaptive problem. In: Cronk L, Chagnon N, Irons W (eds) *Human Behavior and*  
1192 *Adaptation: An Anthropological Perspective*. Aldine, New York, pp 371-395
- 1193 Tillier AM, Arensburg B, Duday H, Vandermeersch B (2001) Brief communication: An early case of  
1194 hydrocephalus: The Middle Paleolithic Qafzeh 12 child (Israel). *Am J Phys Anthropol* 114:166-170
- 1195 Tinbergen N (1963) On aims and methods of ethology. *Z Tierpsychol* 20:410-433
- 1196 Tobias PV (2006) Longevity, death and encephalisation among Plio-Pleistocene hominins. *Int Congr*  
1197 *Ser* 1296:1-15
- 1198 Tomasello M (2014) The ultra-social animal. *European Journal of Social Psychology* 44:187-194
- 1199 Tomasello M, Carpenter M, Call J, Behne T, Moll H (2005) Understanding and sharing intentions: The  
1200 origins of cultural cognition. *Behav Brain Sci* 28:675-735
- 1201 Tomasello M, Hare B, Lehmann H, Call J (2007) Reliance on head versus eyes in the gaze following of  
1202 great apes and human infants: the cooperative eye hypothesis. *J Hum Evol* 52:314-320
- 1203 Trinkaus E, Hillson S, Franciscus RG, Holliday TW (2006) Skeletal and dental paleopathology. In:  
1204 Trinkaus E, Svoboda J (eds) *Early Modern Human Evolution in Central Europe: The People of Dolní*  
1205 *Věstonice and Pavlov*. Oxford University Press, New York, pp 419-458
- 1206 Trinkaus E, Jelinek J (1997) Human remains from the Moravian Gravettian: The Dolni Vestonice 3  
1207 postcrania. *J Hum Evol* 33:33-82
- 1208 Turner SE, Fedigan L, Damon Matthews H, Nakamichi M (2014) Social consequences of disability in a  
1209 nonhuman primate. *J Hum Evol* 68:47-57
- 1210 van Schaik CP, Isler K, Burkart JM (2012) Explaining brain size variation: from social to cultural brain.  
1211 *Trends in Cognitive Sciences* 16:277-284
- 1212 Walker A, Zimmerman MR, Leakey REF (1982) A possible case of hypervitaminosis A in *Homo erectus*.  
1213 *Nature* 296:248-250

- 1214 Weiss RA (2007) Lessons from naked apes and their infections. *J Med Primatol* 36:172-179
- 1215 Whiten A (2000) Primate culture and social learning. *Cognitive Science* 24:477-508
- 1216 WHO (2013) Crimean-Congo Haemorrhagic Fever.
- 1217 WHO (2014a) Ebola Virus Disease.
- 1218 WHO (2014b) Measles.
- 1219 WHO (2015) Scabies.
- 1220 Wilensky U (1999) Netlogo. Center for Connected Learning and Computer-Based Modeling,  
1221 Northwestern University, Evanston, IL
- 1222 Wolfe ND, Dunavan CP, Diamond J (2007) Origins of major human infectious diseases. *Nature*  
1223 447:279-283
- 1224 Woodley MA, Fernandes HBF, Hopkins WD (2015) The more g-loaded, the more heritable, evolvable,  
1225 and phenotypically variable: Homology with humans in chimpanzee cognitive abilities. *Intelligence*  
1226 50:159-163
- 1227 Wu X, Holloway RL, Schepartz LA, Xing S (2011) A new brain endocast of *Homo erectus* from Hulu  
1228 Cave, Nanjing, China. *Am J Phys Anthropol* 145:452-460
- 1229 Zylberberg M, Klasing KC, Hahn TP (2012) House finches (*Carpodacus mexicanus*) balance investment  
1230 in behavioural and immunological defences against pathogens. *Biology Letters* 9:4  
1231  
1232

## Tables

Table 1. Summary data calculated from the hominin dataset presented in Appendix Table A1 of Grove et al. (2012). Values and confidence intervals are medians calculated from the published dataset. To keep our terminology consistent, we refer to community size where Grove et al. (2012) refers to group size.

Genus	Species	Community Size			Population Density (I/km <sup>2</sup> )			Low
		Lower CI	Median	Upper CI	Lower CI	Median	Upper CI	
<i>Homo</i>	Early <i>Homo</i>	43.249	56.276	71.402	0.366	0.584	0.802	51.5
<i>Homo</i>	<i>habilis</i>	46.8415	60.476	76.2795	0.577	0.822	1.068	43.8
<i>Homo</i>	<i>erectus</i>	66.43	83.158	102.406	0.545	0.785	1.025	70.2

<i>Homo heidelbergensis</i>	70.9845	88.389	108.389	0.3	0.514	0.728	94.7
<i>Homo neanderthalensis</i>	72.622	90.266	110.5325	0.196	0.407	0.618	116.
<i>Homo sapiens</i>	78.763	97.292	118.541	0.196	0.407	0.618	127.

Table 2. Means and standard deviations for each disease for the final population size, final disease prevalence, final percent care, final average population intelligence, the net intelligence change between time steps 1 and 100 (Intel Change), the maximum slope for percent care, and the maximum slope for average population intelligence from Model 1 (Care-giving).

Disease	Pop. Size		Prevalence (%)		Percent Care		Intelligence		Intel Change		Slope Care	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Scabies	184.07	0.77	84.78	0.42	1.37	0.11	0.53	0.01	0.03	<0.01	-0.00006	0.0
Measles	133.64	2.02	70.15	0.76	6.74	0.43	0.71	0.01	0.21	0.01	0.00053	0.0
CCHF	120.96	3.47	33.63	1.75	4.73	0.50	0.62	0.01	0.12	0.02	0.00022	0.0
Ebola	157.24	3.25	10.32	0.51	---	---	0.50	0.01	0.00	0.01	---	---

Table 3. One-way ANOVAs showing significant differences across disease types for the final population size, final disease prevalence, final percent care, final average population intelligence, the net intelligence change between time steps 1 and 100, the maximum slope for percent care, and the maximum slope for average population intelligence for Model 1 (Care-giving). All multiple comparisons between disease types were



significant, thus only the smallest mean difference and corresponding p-value are shown per test.

Test	F-statistic	Df	P	Smallest Mean Difference	P
Final Pop. Size	1131.78 <sup>BF</sup>	3, 24.47	<0.001	$\geq 12.68^T$	<0.001
Final Prevalence	11,275.24 <sup>BF</sup>	3, 15.24	<0.001	$\geq 0.15^T$	< 0.001
Final Percent Care	492.03 <sup>BF</sup>	2, 18.61	<0.001	$\geq 0.02^T$	<0.001
Final Intelligence	579.51 <sup>UC</sup>	3, 36	<0.001	$\geq 0.03^B$	<0.001
Intelligence Change	464.463 <sup>BF</sup>	3, 23.13	<0.001	$\geq 0.03^T$	<0.001
Max. Slope Percent Care	377.10 <sup>UC</sup>	2, 27	<0.001	$\geq 0.0003^B$	<0.001
Max. Slope Intelligence	421.732 <sup>BF</sup>	3, 21.61	<0.001	$\geq 0.0002^T$	$\leq 0.03$

<sup>UC</sup> F-statistic, uncorrected

<sup>BF</sup> Brown-Forsythe F-statistic

<sup>B</sup> Bonferroni correction for multiple comparisons

<sup>T</sup> Tamhane's T2 test for multiple comparisons

Table 4. One-sample T-tests on the *Model 1 results* showing that the difference in average population intelligence between the first and 100<sup>th</sup> time steps were significantly different from zero for the scabies-like, measles-like, CCHF-like diseases, but not for the Ebola-like disease. Significant p-values are bolded

Test	T	Df	P	CI: Lower	CI: Upper
Scabies-like	22.18	9	<b>&lt;0.001</b>	0.028	0.033
Measles-like	44.78	9	<b>&lt;0.001</b>	0.196	0.216
CCHF-like	19.36	9	<b>&lt;0.001</b>	0.111	0.137
Ebola-like	-0.824	9	0.431	-0.010	0.005

Table 5. One-sample T-tests on the *Model 2 results* showing that the difference in average population intelligence between the first and 100<sup>th</sup> time steps were significantly different from zero for the CCHF-like and Ebola-like diseases, but not for the scabies-like and measles-like diseases. Significant p-values are bolded.

Test	T	Df	P	CI: Lower	CI: Upper
Scabies-like	-.997	9	0.352	-0.005	0.001
Measles-like	-1.292	9	0.236	-0.025	0.005
CCHF-like	-24.000	9	<b>0.001</b>	-0.160	-0.138
Ebola-like	-58.939	9	<b>0.001</b>	-0.216	-0.200

Table 6. Two-sample T-tests comparing population size, prevalence, and mean intelligence values at the 100<sup>th</sup> time step for each disease under Model 1 (care-giving) versus

Model 2 (avoidance) conditions. When Levene's test indicated that the variances are unequal, we report the T values, degrees of freedom (df), p-values, and confidence intervals calculated without assuming equal variances (Field 2013). Significant p-values are bolded.

Disease	Variable	T	Df	P	CI:	CI:
					Lower	Upper
Scabies-like	Pop. Size	43.178	11.011	<b>0.001</b>	28.833	31.344
	Prevalence	-49.675	18	<b>0.001</b>	-0.105	-0.096
	Intelligence	7.786	18	<b>0.001</b>	0.031	0.052
Measles-like	Pop. Size	9.669	18	<b>0.001</b>	9.621	14.569
	Prevalence	-3.000	18	<b>0.016</b>	-0.029	-0.007
	Intelligence	30.699	11.148	<b>0.001</b>	0.205	0.233
CCHF-like	Pop. Size	-3.165	18	<b>0.003</b>	-5.906	-1.296
	Prevalence	0.740	18	0.464	-0.007	0.015
	Intelligence	37.944	18	<b>0.001</b>	0.254	0.282
Ebola-like	Pop. Size	-0.024	14.171	0.982	-3.696	3.923
	Prevalence	0.305	18	0.748	-0.004	0.005
	Intelligence	46.049	18	<b>0.001</b>	0.200	0.218

Table 7. Mixed-model analyses run on the Model 1 (care-giving) results examining the effects of prevalence, population size and the interaction between the two on intelligence changes for each disease.  $r^2_m$  measures how much variation in mean intelligence can be explained by the fixed effects (time+prevalence\*population size).

$\beta$  values are standardized regression coefficients. SE is the standard error and df is the degrees of freedom.


Disease	Analysis	$r^2_m$ *	Variable	B	SE	df	t	
Scabies-like*	Prevalence	0.468	Intercept	-0.002	0.034	888	-0.055	0
			Time	-1.084	0.086	888	12.641	<0
			Prevalence	0.460	0.085	888	5.411	<0
Measles-like	Prevalence	0.565	Intercept	-0.065	0.075	946	-0.871	0
			Time	-0.585	0.076	946	-7.650	<0
			Population Size	0.291	0.063	946	4.590	<0
			Prevalence	0.431	0.046	946	9.276	<0
			Population Size*Prevalence	-0.143	0.021	946	-6.713	<0
CCHF-like	Prevalence	0.146	Intercept	0.039	0.050	946	0.785	0
			Time	-0.400	0.051	946	-7.848	<0
			Population Size	0.052	0.051	946	1.014	0
			Prevalence	-0.104	0.052	946	-2.023	0
			Population Size*Prevalence	0.060	0.020	946	3.023	0
Ebola-like	Prevalence	0.001	Intercept	0.008	0.039	946	0.218	0
			Time	-0.010	0.039	946	-0.247	0
			Population Size	-0.043	0.049	946	-0.873	0
			Prevalence	0.002	0.073	946	0.031	0
			Population Size*Prevalence	0.013	0.022	946	0.571	0


\* $r^2_c$  values were the same as  $r^2_m$ .  $r^2_c$  measures how much variation is explained by the whole

model (including the random effect of simulation run). That the two measures were the same

indicates that there were no systematic differences between runs of a given disease.

Figure 1. Changes over time in disease prevalence (A), percentage of diseased individuals who received care (B), and average population intelligence (C). For each disease the 10 average runs have been averaged within each time step. The Ebola-like, CCHF-like, measles-like, and scabies-like diseases are shown in red circles, green squares, black Xs, and blue triangles, respectively. Approximately every fourth time step is shown. Error bars are  $\pm$  two standard deviations. Fig. 1B does not show the Ebola-like disease because no care was given.











Figure 2. Changes in population size over time produced by Model 1 (Care-giving) and Model 2 (Avoidance only) in the scabies-like (A), measles-like (B), CCHF-like (C), and Ebola-like (D) diseases. Models 1 and 2 are shown in red circles and blue triangles, respectively. Only even numbered time steps are shown. Error bars are  $\pm$  two standard deviations.















Figure 3. Changes in prevalence over time produced by Model 1 (Care-giving) and Model 2 (Avoidance only) in the scabies-like (A), measles-like (B), CCHF-like (C), and Ebola-like (D) diseases. Models 1 and 2 are shown in red circles and blue triangles, respectively. Only even numbered time steps are shown. Error bars are  $\pm$  two standard deviations.













Figure 4. Changes in average population intelligence over time produced by Model 1 (Care-giving) and Model 2 (Avoidance only) in the scabies-like (A), measles-like (B), CCHF-like (C), and Ebola-like (D) diseases. Models 1 and 2 are shown in red circles and blue triangles, respectively. Only even numbered time steps are shown. Error bars are +/- two standard deviations.












Figure 5. Graphs showing the results of the analyses exploring the effects of prevalence on the change in intelligence for the scabies-like disease. Change in intelligence was calculated as the mean intelligence in a given time step minus the mean intelligence in the previous time step. (A) Change in intelligence is negatively correlated with time and (B) positively correlated with prevalence (Table 7).




Figure 6. Graphs showing the results of the analyses exploring the effects of prevalence, population size, and their interactions on the change in intelligence for the measles-like disease. Change in intelligence was calculated as the mean intelligence in a given time step minus the mean intelligence in the previous time step. (A) Change in intelligence is negatively correlated with time (Table 7). (B) Interaction effects between population size and prevalence (“Prev”). Population size is on the X axis with data points represented by the small black lines. The difference in intelligence is shown on the Y axis. The prevalences shown represent the range of prevalences experienced by the population (see Figure 1A). The greatest positive selection on intelligence occurred when prevalence and population size are high. Population size has a large effect when prevalence is low (left panel of B) and a small effect when prevalence is high (right panel of B).




Figure 7. Graphs showing the results of the analyses exploring the effects of prevalence, population size, and their interactions on the change in intelligence for the CCHF-like disease. Change in intelligence was calculated as the mean intelligence in a given time step minus the mean intelligence in the previous time step. (A) Change in intelligence is negatively correlated with time (Table 7). (B) Interaction effects between population size and prevalence. Population size is on the X axis with data points represented by the small black lines. The difference in intelligence is shown on the Y axis. The prevalences shown represent the range of prevalences experienced by the population (see Figure 1A). The greatest increases in average population intelligence occurred at low population sizes and low prevalences (B, left panel) and at high population sizes and high prevalences (B, right panel).




Figure 8. Graphs showing the results of the analyses exploring the effects of prevalence, population size, and their interactions on change in intelligence for the Ebola-like disease. Change in intelligence was calculated as the mean intelligence in a given time step minus the mean intelligence in the previous time step. (A) No significant change in intelligence over time. (B) Potential interaction effects between population size and prevalence. Population size is on the X axis with data points represented by the small black lines. The difference in intelligence is shown on the Y axis. The prevalences shown represent the range of prevalences experienced by the population (see Figure 1A). Because intelligence does not change over time, there are no significant correlations with prevalence, population size or the interaction of the two (Table 7).

